

NEA OYH META TON KTIPIAKO EKSTYXONISMO

ΕΚΘΕΣΙΑΚΟ ΚΑΙ ΕΜΠΟΡΙΚΟ
ΚΕΝΤΡΟ "LINEA"

SYNOPSIS

NEPHTHALYNOYL: XANTHENEAS & AHA. EPFON

[illegible]

AGHINA 8 ZENT 48 pios 2000

10/11/17

[Signature]
C. Adams

1. Protein

10

10

[illegible]

УПРАВЛЕНИЕ
ИСПИТАНИЯ

ΟΤΑΕΙΑΣ & ΔΗΜΟΣΙ

ΠΟΛΥΕΘΔΟΜΙΚΟΥ ΣΧΕΔΙΑΣ

ΠΑΡΑΔΟΧΙΑΚΗΝ ΔΙΚΙΣΜΟΝ

1

ΔΗΜΟΣ ΠΕΙΡΑΙΑ

1

ΕΛΛΗΝΙΚΗ ΔΗΜΟΚΡΑΤΙΑ
ΥΠΟΥΡΓΕΙΟ ΠΑΙΔΕΙΑΣ, ΕΡΕΥΝΑΣ ΚΑΙ ΘΡΗΣΚΕΥΜΑΤΩΝ
ΙΝΣΤΙΤΟΥΤΟ ΤΕΧΝΟΛΟΓΙΑΣ ΥΠΟΛΟΓΙΣΤΩΝ ΚΑΙ ΕΚΔΟΣΕΩΝ ΔΙΔΑΚΤΙΚΩΝ ΒΙΒΛΙΩΝ (ΙΤΥΥΔΕΚ)

1. *Chlorophyll a* and *Chlorophyll b* were determined by the method of Lichtenthaler and Whistler (1973). The total chlorophyll content was determined by the method of Arar and Cook (1980). The carotenoid content was determined by the method of Lichtenthaler and Whistler (1973). The total carotenoid content was determined by the method of Arar and Cook (1980). The total protein content was determined by the method of Lowry et al. (1951). The total lipid content was determined by the method of Bligh and Dyer (1959). The total carbohydrate content was determined by the method of Dubois and Gilles (1950). The total nucleic acid content was determined by the method of Burton (1956). The total ash content was determined by the method of AOAC (1990). The total fiber content was determined by the method of AOAC (1990). The total organic acid content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenol content was determined by the method of AOAC (1990). The total terpenoid content was determined by the method of AOAC (1990). The total steroid content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenol content was determined by the method of AOAC (1990). The total terpenoid content was determined by the method of AOAC (1990). The total steroid content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990).

Τροπικότητα ειδικών ορών δομής

[illegible]

Καραγόλη - Δημητρίου, Μιχαήλ

—

Printed in the U.S.A. by MAPT 102 2000

1

0
1
2
3
4
5
6
7
8
9
A
B
C
D
E
F
G
H
I
J
K
L
M
N
O
P
Q
R
S
T
U
V
W
X
Y
Z

OUT QUADRA LARVAE AND MILK

Ad. Ph.

111

10

7